

Simulation of $\gamma p \rightarrow K^+ K^+ \Xi^- (1530)$

Reaction

$$\gamma p \rightarrow K^+ K^+ \Xi^-(1530),$$

Reaction

$$\gamma p \rightarrow K^+ K^+ \bar{E}(1530),$$

where

$$\bar{E}(1530) \rightarrow \bar{E} \pi^0,$$

Reaction

$$\gamma p \rightarrow K^+ K^+ \bar{E}(1530),$$

where

$$\bar{E}(1530) \rightarrow \bar{E} \pi^0,$$

$$\bar{E} \rightarrow \Lambda \pi,$$

Reaction

$$\gamma p \rightarrow K^+ K^+ \bar{E}(1530),$$

where

$$\bar{E}(1530) \rightarrow \bar{E} \pi^0,$$

$$\bar{E} \rightarrow \Lambda \pi,$$

and

$$\Lambda \rightarrow p \pi^-$$

E^* Generator Refinement

- Started with generator for general reaction $\gamma p \rightarrow K^+ K^+ E^- \pi^0$ and modified from there
- Taking the initial reaction as $\gamma p \rightarrow K Y^*$
- Mandelstam variables have relationship:
 - $s+t+u = m_\gamma^2 + m_p^2 + m_K^2 + m_{Y^*}^2$
- We can lock down the kinematics of the initial reaction by specifying s , t and m_{Y^*}
- Started with Mandelstam s and t

Ξ^* Comparison of Reconstructed MC to Actual Data

For the general reaction:


- 1st: Set t -slope (parameter b in $Ae^{-b|t|}$) to $1.138/\text{GeV}^2$ assuming $\gamma p \rightarrow K_{\text{fast}} Y^*$
- 2nd: Shaped mass[Y^*] and mass[Ξ^*]
- 3rd: Shaping K_{slow} distribution assuming $Y^* \rightarrow K_{\text{slow}} \Xi^*$

For $\gamma p \rightarrow K^+ K^+ \Xi^-(1530)$


- Start with general reaction and modify each step as required

E^* Comparison of Reconstructed MC to Actual Data

General reaction




$\gamma p \rightarrow K^+ K^+ E^-(1530)$






E^* Comparison of Reconstructed MC to Actual Data

General reaction




$\gamma p \rightarrow K^+ K^+ E^-(1530)$




Ξ^* Comparison of Reconstructed MC to Actual Data

General reaction




$\gamma p \rightarrow K^+ K^+ \Xi^-(1530)$






E^* Comparison of Reconstructed MC to Actual Data

General reaction




$\gamma p \rightarrow K^+ K^+ \Xi^-(1530)$




Ξ^* Comparison of Reconstructed MC to Actual Data

General reaction




$\gamma p \rightarrow K^+ K^+ \Xi^-(1530)$




Y^* Comparison of Reconstructed MC to Actual Data

General reaction



$\gamma p \rightarrow K^+ K^+ Y^*(1530)$




Resonances in the Y^* ?




Will explore more in later slides

E^* Comparison of Reconstructed MC to Actual Data

General reaction




$\gamma p \rightarrow K^+ K^+ E^-(1530)$




E^* Comparison of Reconstructed MC to Actual Data

General reaction




$\gamma p \rightarrow K^+ K^+ E^-(1530)$




E^* Comparison of Reconstructed MC to Actual Data

General reaction



$\gamma p \rightarrow K^+ K^+ E^-(1530)$



Not good ☹


Y^* bumps compared to PDG Σ bumps

Y^* bumps compared to PDG Σ bumps

$\gamma p \rightarrow K^+ Y^*$, where $Y^* \rightarrow K^+ \Xi^-(1530)$ and $\Xi^-(1530) \rightarrow \Xi^- \pi^0$

Y^* bumps compared to PDG Σ bumps

$\gamma p \rightarrow K^+ Y^*$, where $Y^* \rightarrow K^+ \Xi^-(1530)$ and $\Xi^-(1530) \rightarrow \Xi^- \pi^0$




First bump:

- Center = 2610 +/- 81 MeV
- FWHM = 353 +/- 270 MeV

Y^* bumps compared to PDG Σ bumps

$$\gamma p \rightarrow K^+ Y^*, \text{ where } Y^* \rightarrow K^+ \Xi^-(1530) \text{ and } \Xi^-(1530) \rightarrow \Xi^- \pi^0$$

Citation: S. Navas *et al.* (Particle Data Group), Phys. Rev. D **110**, 030001 (2024) and 2025 update



$\Sigma(2620)$ Bumps

$I(J^P) = 1(?^?)$ Status: *

OMITTED FROM SUMMARY TABLE

$\Sigma(2620)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|-----------------------------------|
| ≈ 2620 OUR ESTIMATE | | | |
| 2542 ± 22 | DIBIANCA | 75 | DBC $K^- N \rightarrow \Xi K \pi$ |
| 2620 ± 15 | ABRAMS | 70 | CNTR $K^- p, K^- d$ total |

$\Sigma(2620)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|--------------|-------------|------|-----------------------------------|
| 221 ± 81 | DIBIANCA | 75 | DBC $K^- N \rightarrow \Xi K \pi$ |
| 175 | ABRAMS | 70 | CNTR $K^- p, K^- d$ total |


First bump:

- Center = 2610 ± 81 MeV
- FWHM = 353 ± 270 MeV

Y^* bumps compared to PDG Σ bumps

$\gamma p \rightarrow K^+ Y^*$, where $Y^* \rightarrow K^+ \Xi^-(1530)$ and $\Xi^-(1530) \rightarrow \Xi^- \pi^0$

Citation: S. Navas *et al.* (Particle Data Group), Phys. Rev. D **110**, 030001 (2024) and 2025 update



| $\Sigma(2620)$ Bumps | | $I(J^P) = 1(?)$ | | Status: * |
|--|-------------|-----------------|---------|-------------------------------|
| OMITTED FROM SUMMARY TABLE | | | | |
| $\Sigma(2620)$ MASS | | | | |
| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT | |
| ≈ 2620 OUR ESTIMATE | | | | |
| 2542 ± 22 | DIBIANCA | 75 | DBC | $K^- N \rightarrow \Xi K \pi$ |
| 2620 ± 15 | ABRAMS | 70 | CNTR | $K^- p, K^- d$ total |
| $\Sigma(2620)$ WIDTH | | | | |
| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT | |
| 221 ± 81 | DIBIANCA | 75 | DBC | $K^- N \rightarrow \Xi K \pi$ |
| 175 | ABRAMS | 70 | CNTR | $K^- p, K^- d$ total |

First bump:


- Center = 2610 ± 81 MeV
- FWHM = 353 ± 270 MeV

Similar final state to
 $\gamma p \rightarrow K^+ K^+ \Xi^- \pi^0$

Y^* bumps compared to PDG Σ bumps

$\gamma p \rightarrow K^+ Y^*$, where $Y^* \rightarrow K^+ \Xi^-(1530)$ and $\Xi^-(1530) \rightarrow \Xi^- \pi^0$

Citation: S. Navas *et al.* (Particle Data Group), Phys. Rev. D **110**, 030001 (2024) and 2025 update



$\Sigma(2620)$ Bumps

$I(J^P) = 1(?^?)$ Status: *

OMITTED FROM SUMMARY TABLE

$\Sigma(2620)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|-----------------------------------|
| ≈ 2620 OUR ESTIMATE | | | |
| 2542 ± 22 | DIBIANCA | 75 | DBC $K^- N \rightarrow \Xi K \pi$ |
| 2620 ± 15 | ABRAMS | 70 | CNTR $K^- p, K^- d$ total |

$\Sigma(2620)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|--------------|-------------|------|-----------------------------------|
| 221 ± 81 | DIBIANCA | 75 | DBC $K^- N \rightarrow \Xi K \pi$ |
| 175 | ABRAMS | 70 | CNTR $K^- p, K^- d$ total |

First bump:

- Center = 2610 ± 81 MeV
- FWHM = 353 ± 270 MeV


Second bump:

- Center = 3093 ± 55 MeV
- FWHM = 240 ± 167

Y^* bumps compared to PDG Σ bumps

$\gamma p \rightarrow K^+ Y^*$, where $Y^* \rightarrow K^+ \Xi^-(1530)$ and $\Xi^-(1530) \rightarrow \Xi^- \pi^0$

Citation: S. Navas *et al.* (Particle Data Group), Phys. Rev. D **110**, 030001 (2024) and 2025 update



First bump:

- Center = 2610 +/- 81 MeV
- FWHM = 353 +/- 270 MeV

Second bump:

- Center = 3093 +/- 55 MeV
- FWHM = 240 +/- 167

$\Sigma(2620)$ Bumps

$I(J^P) = 1(?^?)$ Status: *

OMITTED FROM SUMMARY TABLE

$\Sigma(2620)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|-----------------------------------|
| ≈ 2620 OUR ESTIMATE | | | |
| 2542 ± 22 | DIBIANCA | 75 | DBC $K^- N \rightarrow \Xi K \pi$ |
| 2620 ± 15 | ABRAMS | 70 | CNTR $K^- p, K^- d$ total |

$\Sigma(2620)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|-----------------------------------|
| 221 ± 81 | DIBIANCA | 75 | DBC $K^- N \rightarrow \Xi K \pi$ |
| 175 | ABRAMS | 70 | CNTR $K^- p, K^- d$ total |

Citation: S. Navas *et al.* (Particle Data Group), Phys. Rev. D **110**, 030001 (2024) and 2025 update

$\Sigma(3000)$ Bumps

$I(J^P) = 1(?^?)$ Status: *

OMITTED FROM SUMMARY TABLE

Seen as an enhancement in $\Lambda \pi$ and $\bar{K} N$ invariant mass spectra and in the missing mass of neutrals recoiling against a K^0 .

$\Sigma(3000)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | CHG | COMMENT |
|-----------------------------|-------------|------|-----|------------------------|
| ≈ 3000 OUR ESTIMATE | | | | |
| 3000 | EHRlich | 66 | HBC | 0 $\pi^- p$ 7.91 GeV/c |


$\Sigma(3000)$ DECAY MODES

| | Mode |
|------------|---------------|
| Γ_1 | $N \bar{K}$ |
| Γ_2 | $\Lambda \pi$ |

Y^* bumps compared to PDG Σ bumps

$\gamma p \rightarrow K^+ Y^*$, where $Y^* \rightarrow K^+ \Xi^-(1530)$ and $\Xi^-(1530) \rightarrow \Xi^- \pi^0$

Citation: S. Navas *et al.* (Particle Data Group), Phys. Rev. D **110**, 030001 (2024) and 2025 update



First bump:

- Center = 2610 +/- 81 MeV
- FWHM = 353 +/- 270 MeV

Second bump:

- Center = 3093 +/- 55 MeV
- FWHM = 240 +/- 167

Interesting, but needs more statistics

$\Sigma(2620)$ Bumps

$I(J^P) = 1(?^?)$ Status: *

OMITTED FROM SUMMARY TABLE

$\Sigma(2620)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-----------------------------|-------------|------|-----------------------------------|
| ≈ 2620 OUR ESTIMATE | | | |
| 2542 ± 22 | DIBIANCA | 75 | DBC $K^- N \rightarrow \Xi K \pi$ |
| 2620 ± 15 | ABRAMS | 70 | CNTR $K^- p, K^- d$ total |

$\Sigma(2620)$ WIDTH

| VALUE (MeV) | DOCUMENT ID | TECN | COMMENT |
|-------------|-------------|------|-----------------------------------|
| 221 ± 81 | DIBIANCA | 75 | DBC $K^- N \rightarrow \Xi K \pi$ |
| 175 | ABRAMS | 70 | CNTR $K^- p, K^- d$ total |

Citation: S. Navas *et al.* (Particle Data Group), Phys. Rev. D **110**, 030001 (2024) and 2025 update

$\Sigma(3000)$ Bumps

$I(J^P) = 1(?^?)$ Status: *

OMITTED FROM SUMMARY TABLE

Seen as an enhancement in $\Lambda \pi$ and $\bar{K} N$ invariant mass spectra and in the missing mass of neutrals recoiling against a K^0 .

$\Sigma(3000)$ MASS

| VALUE (MeV) | DOCUMENT ID | TECN | CHG | COMMENT |
|-----------------------------|-------------|------|-----|------------------------|
| ≈ 3000 OUR ESTIMATE | | | | |
| 3000 | EHRlich | 66 | HBC | 0 $\pi^- p$ 7.91 GeV/c |

$\Sigma(3000)$ DECAY MODES

| | Mode |
|------------|---------------|
| Γ_1 | $N \bar{K}$ |
| Γ_2 | $\Lambda \pi$ |

Title

